

Unehäirete ravijuhendi kliiniline küsimus 16:

Kas kõigil liigunisuse kahtlusega patsientidel tuleb kasutada uneaegse hingamishäire küsimustikku vs mitte?

28.03.2018

PICO	Otsingustrateegia tulemus	
<p>P adults with excessive daytime sleepiness or with excessive sleepiness or with daytime sleepiness or with excessive daytime somnolence</p> <p>I sleep-related breathing disorder questionnaires or sleep-related breathing disorder scales or sleep-related breathing disorder inventory or respiratory sleep-related breathing disorders screening tools or respiratory sleep-related breathing disorders screening questionnaires or sleep apnea questionnaires or apnea screening tools or Epworth sleepiness scale</p> <p>C not using questionnaire or not screening or or interviewing</p> <p>O sleep apnea or or sleep related breathing disorder or breathing-related sleep disorder or hypersomnolence or hypersomnia</p>	<p>Kokku 866 inglise keelset kokkuvõtet</p>	<ol style="list-style-type: none"> 1. Liigunisuus, OUA, mootorsõidukijuhid 2. Narkolepsia 3. Liigunisuus, insomnia MS, Parkinsonism 4. Liigunisuus, bariaatriline kirurgia 5. CPAP 6. jne <p>+ uuringud, mis kattusid kliiniliste küsimustega:</p> <ol style="list-style-type: none"> 1. Kas kõigil uneaegse hingamishäire kahtlusega patsientidel tuleb kasutada uneaegsete hingamishäirete küsimustikku vs mitte? 2. Kas kõigil uneaegse hingamishäire kahtlusega patsientidel tuleb kasutada liigunisuse küsimustikku vs mitte?
<p>P adults with excessive daytime sleepiness or with excessive sleepiness or with daytime sleepiness or with excessive daytime somnolence</p> <p>I STOP BANG questionnaires, STOP questionnaire</p>	<p>Tulemuseks 25 kirjet. Pärast ajalise ja keelepiirangu lisamist (publikatsioonid alates 2003. aastast; keel: English) jäi nimekirja 20 kirjet</p>	<p>NB! Sobilikke ei leidunud</p>